

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/13, 31/445, 31/55, 31/495, 31/35, 31/415		A2	(11) International Publication Number: WO 95/21611
			(43) International Publication Date: 17 August 1995 (17.08.95)
(21) International Application Number: PCT/US95/01392			(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).
(22) International Filing Date: 9 February 1995 (09.02.95)			
(30) Priority Data: 08/193,597 9 February 1994 (09.02.94) US			
(71) Applicant: KOS PHARMACEUTICALS, INC. [US/US]; Suite 2502, 1001 South Bayshore Drive, Miami, FL 33131 (US).			
(72) Inventor: THEOHARIDES, Theoharis, C.; 14 Parkman Street, Brookline, MA 02146 (US).			
(74) Agents: MELOY, Sybil et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).			Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: COMPOSITIONS COMPRISING A MAST CELL DEGRANULATION-BLOCKING AGENT FOR TREATMENT OF ENDOGENOUS, PAINFUL GASTROINTESTINAL CONDITIONS OF NON-INFLAMMATORY, NON-ULCERATIVE ORIGIN			
(57) Abstract <p>A method for treating endogenous, painful gastrointestinal conditions of non-inflammatory, non-ulcerative origin, such as abdominal migraine and irritable bowel syndrome, entails administering a pharmacologically effective amount of a mast cell degranulation-blocking agent, e.g. a naturally occurring polyamine, a heterocyclic histamine-1 receptor antagonist, a histamine-3 receptor agonist or an anti-female sex hormone.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

COMPOSITIONS COMPRISING A MAST CELL DEGRANULATION-BLOCKING
AGENT FOR TREATMENT OF ENDOGENOUS, PAINFUL GASTROINTESTINAL
CONDITIONS OF NON-INFLAMMATORY, NON-ULCERATIVE ORIGIN

5

BACKGROUND OF THE INVENTION

The present invention relates to treating endogenous, painful gastrointestinal conditions, such as abdominal migraine and irritable bowel syndrome, which involve neither inflammation nor ulcers. "Endogenous" in this context denotes a condition that is not attributed to an exogenous casual factor such as a food allergy, a bacterial or viral infection, a parasitic infestation, a drug reaction or trauma. More specifically, the present invention relates to treating a patient suffering from such a condition with a pharmaceutically effective amount of a mast cell degranulation-blocking agent.

The phrase "abdominal pain" indicates pain associated with the gastrointestinal tract. Functional abdominal pain usually is classified either as dyspepsia not associated with ulcers or as irritable bowel syndrome.

Irritable bowel syndrome (IBS), also known as "spastic colon" and "mucous colitis," is an intestinal motility disorder and ranks among the most common pathological conditions of the intestine. IBS is characterized by periodic or chronic bowel symptoms which include abdominal pain, diarrhea, constipation, a sense of incomplete evacuation, bloating, and excess gas sensation. This disorder seems to afflict type A (highly driven, perfectionist) personalities predominantly, and is two to five times more prevalent in women (20 times higher in Jewish women) than in men. It affects about 3 percent of the population. Drossman, *Hospital Practice* 93: 95-108 (1988).

Pain and flatulence are the most prominent symptoms in patients suffering from IBS. The pain is usually situated in the left lower quadrant or suprapubically. It may be worse just before defecation and may lessen slowly afterward. Certain foods may precipitate the symptoms. When diarrhea is the main complaint, the stool often is watery but not bloody. The diarrhea frequently is worst in the morning and improves during the day.

Some patients complain of pain in the left upper quadrant that is often brought on by meals and is very consistent in nature. In these patients, plain abdominal X-rays demonstrate air in the splenic flexure. Although this variant of IBS has been called "splenic flexure syndrome," there is no evidence that its pathogenesis differs from that of other manifestations of the disorder.

Many patients relate an exacerbation of symptoms to episodes of emotional stress. Fear of underlying cancer is common. In children of school age, IBS presents primarily with pain. Pain is usually periumbilical or in the left lower quadrant and is cramping in nature.

Many patients also experience abdominal pain that is endogenous but that is associated with nausea, bowel peristalsis and flatulence without diarrhea or constipation, the symptoms commonly seen with IBS. Such abdominal pain also can be associated with classic symptoms of migraine, such as one-sided headache with possible involvement of one eye and visual disturbances in that eye. Axon et al., *J. Clin. Gastroenterol.* 13: 615-16 (1991). By the same token, about 30-50% of patients diagnosed with IBS have frequent headaches, as compared to less than 15% in the control population. Watson et al., *Can. Med. Assoc. J.* 118: 387-98 (1978); Whorwell et al., *Gut* 27: 37-40 (1986). Finally, abdominal pain and associated symptoms can occur in the absence of a headache and still be considered a "migraine equivalent," especially in children. Lundbert, *Headache* 15: 122-25 (1975).

It is appropriate, therefore, to consider the pathophysiology and therapy of the category of endogenous, painful gastrointestinal conditions of non-inflammatory, non-ulcerative origin, including but not limited to abdominal migraine and IBS.

There is no effective therapy for IBS or related, endogenous gastrointestinal conditions. Thus, while various compounds have been described as useful in treating IBS, including 9H-fluorenyl-substituted amino acid derivatives (U.S. patent No. 5,079,260), benzodiazepine derivatives (U.S. patent No. 4,970,207), certain substituted sulfonamides (European application No. 0 404 440), and mu opiate receptor antagonists or "blockers" (U.S. patent No. 4,684,620), none of the disclosed treatment approaches in fact have proven successful. See, for example, *J. Gastroenterol.* 95: 232-41 (1988).

Similarly, no effective remedy exists for abdominal migraine. Development of a treatment has been hindered in part by the fact that the condition often presents without headache and any of the prodromal symptoms (nausea, vomiting, photophobia, sonophobia) typically associated with ordinary migraine.

For example, agents that prevent vasodilation, such as the β -blocker propranolol, seem not to work, since the abdominal pain appears not to be associated with the vasculature, in contrast to the migraine headache. By the same token, none of the drugs used prophylactically or acutely (symptomatically) for the treatment of migraine headaches would be expected to work, since the prophylactic agents prevent vasodilation while the acute agents constrict dilated vessels. Moreover, drugs effective against migraine headaches must cross the blood-brain barrier, while a drug effective against abdominal migraine would be concentrated preferentially in the intestine. Also, intracranial mast cells which have been associated with migraine headaches, see Theoharides, *Life Sciences* 46: 607-17 (1980), and U.S.

patent No. 5,250,529, differ from gastrointestinal mast cells. For instance, the former but not the latter are inhibited by disodium cromoglycate. Pierce et al., *J. Immunol.* 128: 2481-86 (1982); Labracht-Hall et al.,
5 *Neuroscience* 39: 199-207 (1990).

SUMMARY OF THE INVENTION

It is an object of the present invention, therefore, to provide an effective treatment for the above-described category of endogenous gastrointestinal conditions which
10 avoids the lack of success and other problems associated with past efforts to this end.

In accomplishing this objective and others, there has been provided, in accordance with one aspect of the present invention, a method for treating endogenous,
15 painful gastrointestinal conditions of non-inflammatory, non-ulcerative origin, which method comprises administering to a patient a pharmacologically effective amount of a mast cell-degranulation blocking agent.

In a preferred embodiment, the mast cell-degranulation blocking agent is selected from the group consisting of naturally occurring polyamines, heterocyclic histamine-1 receptor antagonists, histamine-3 receptor agonists, and agents that counteract the influence of female sex hormones (female sex hormone
20 antagonists or inhibitors, referred to collectively as "anti-female sex hormones"). In other preferred embodiments the mast cell-degranulation blocking agent is spermine or spermidine, hydroxyzine or ketotifen, α,β -difluoro-N^α-fluoromethyl histamine, tamoxifen or a
25 gonadotropin-releasing hormone (leutinizing hormone releasing hormone) analogue. In yet other preferred embodiments, the mast cell-degranulation blocking agent is N^α-dimethyl histamine, R^α-methyl histamine, α,β -dimethyl-N^α(dimethyl) histamine, α,β -difluoro-N^α(dimethyl)
30 histamine, α -methyl, β -fluoro-N^α(fluoromethyl) histamine,
35

clomiphene, mifepristone, tamoxifen, and a GnRH analogue such as LUPRON.

Other objects, features and advantages of the present invention will become apparent to those skilled in the art from the following detailed description. It is to be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the present invention, are given by way of illustration and not limitation. Many changes and modifications within the scope of the present invention may be made without departing from the spirit thereof, and the invention includes all such modifications.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Mast cells are normal components of the connective and mucosal tissues and play an important role in allergy. These cells are mostly localized in the gastrointestinal mucosa, skin and lung. Mast cells are so located, it is believed, because these tissues are the main entry points for infective organisms and allergens, the chemicals that trigger the body's immune response.

Recent evidence indicates that mast cells in the mucosa of the small intestine differ from mast cells in other tissues, especially connective tissues as well as brain. These mucosal mast cells also have been shown to be in close proximity to nerve endings. See Stead et al., *J. Gastroenterol.* 97: 575-85 (1989).

Every mast cell contains up to 500 secretory granules, each storing more than 20 potent biological compounds. Mast cells secrete the contents of these granules, i.e., degranulate, when triggered by various specific and non-specific mechanisms. Accordingly, "degranulation" is used in this description to denote the release of any or all mediators from any or all mast cell secretory granules, whether in parallel, differentially or selectively.

The degranulation of mast cells in response to various agents is usually a biological consequence of the activation of one or more receptors located on the surface of the mast cell. The best known of these are the immunoglobulin E (IgE) receptors involved in allergic reactions, to date the most fully characterized pathological processes that involve mast cells. In this context IgE binds strongly to mast cells through its Fc receptor. When mast cell-bound IgE reacts with an antigen, the latter bridges two or more IgE molecules and causes mast cell degranulation with subsequent release of mediators, either stored or synthesized during mast cell degranulation.

There also is evidence that neurotransmitters such as acetylcholine and neuropeptides such as substance P, all being molecules that are released from neurons in the central (brain) and peripheral (gastrointestinal) nervous systems, may augment or trigger mast cell degranulation primarily through specific receptors, for example, in response to stress. In particular, research has shown that direct nerve stimulation causes mast cell activation and release of relevant chemical agents. See, for instance, Dimitriadou et al., *Neuroscience* 44: 97-112 (1991); Theoharides, *Life Sciences* 46: 607-17 (1990). Female sex hormones such as estradiol and progesterone likewise may augment or trigger mast cell degranulation. See Vliagoftis et al., *Int'l Arch. Allergy Immunol.* 98: 398-409 (1992); 93: 113-19 (1990). Thus, it has been shown that rat basophilic leukemia (RBL) cells, which mimic mucosal mast cells, bind progesterone. Kouretas et al., *Clinical Res.* 39: 320 (1991).

Other triggers of mast cell degranulation include viral and bacterial toxins, drugs such as aspirin, morphine and curare, contrast media used in radiology, extreme heat, cold, solar radiation and hyperosmotic media. It is deemed important, therefore, to block the mast cell degranulation which all of these stimuli

elicit, rather than to antagonize specific receptors, such as opioid receptors as taught in U.S. patent No. 4,684,620.

5 The compounds released by the mast cells following degranulation are known to cause many biological processes which are part of the overall response of the body to invasion by infective organisms and allergens. Examples of such processes are vasoconstriction or vasodilation, leukocyte chemotaxis, and pain.

10 Compounds released by mast cell degranulation which may be associated with irritable bowel syndrome include bradykinin, histamine, leukotrienes, prostaglandins, serotonin, tumor necrosis factor and vasoactive intestinal peptide. Interleukins and proteolytic
15 enzymes, which can affect intestinal function, can augment pain or cause pain directly.

Histamine and other mediators bind to specific receptors on the surface of endothelial cells on vessels or on peripheral sensory neurons which become activated,
20 directly or indirectly (through vessel tone), to transmit pain stimuli. Vasodilation and chemoattraction permit lymphocytes to leave the circulation and enter the tissue, where they cause additional mast cell degranulation and other responses. The process of
25 degranulation continues, eventually involving many mast cells.

Mast cell degranulation, especially in response to neurohormonal triggers, now has been discovered to contribute to the symptoms experienced by patients
30 suffering from abdominal migraine, irritable bowel syndrome and other endogenous, painful gastrointestinal conditions of non-inflammatory, non-ulcerative origin. It also has been found an effective treatment for such conditions entails the administration of a mast cell
35 degranulation-blocking agent, i.e., a compound that counteracts mast cell degranulation-inducing stimuli, either by binding to specific receptors or by inactivating crucial steps in the degranulation process.

The aforementioned correlation between mucosal mast cells and stimuli associated with IBS and related, endogenous conditions has not been documented heretofore. For instance, histologic sections of colonic tissue
5 biopsied from a patient with irritable bowel syndrome were stained with monoclonal anti-tryptase antibody to identify tryptase-immunoreactive mast cells, and other sections were stained with polyclonal anti-SP antibody to reveal nerve fibers positive for substance P, a
10 nociceptive neuropeptide capable of degranulating mast cells. Microscopic examination of the sections revealed numerous mast cells, identifiable by staining for tryptase, a proteolytic enzyme which is specific for the mast cells, and numerous nerve fibers stained for their
15 content of substance P. Particularly noteworthy in this context was the close anatomic association observed to exist between mucosal mast cells and substance P-containing nerve fibers.

For the purposes of this description, the range of
20 "mast cell degranulation-inducing stimuli" is illustrated by: (i) immune molecules, for example, immunoglobulin (IgE), anaphylatoxins such as C5a, and cytokines such as tumor necrosis factor; (ii) neural substances, including neurotransmitters such as acetylcholine and neuropeptides
25 such as substance P, calcitonin gene related peptide and vasoactive intestinal peptide; (iii) hormones, such as estrogens and progestins; and (iv) any other molecules that may be liberated within the body during endogenous conditions such as stress. Particularly preferred mast
30 cell degranulation-blocking agents are: (a) natural polyamines; (b) certain heterocyclic histamine-1 receptor antagonists; (c) histamine-3 receptor agonists and (d) anti-female sex hormones.

Among the preferred polyamines are spermine and
35 spermidine, as well as metabolites, including acetylated metabolites thereof and oxidized metabolites, of these two compounds. See Vliagoftis et al., *Biochem.*

Pharmacol. 43: 2237-45 (1992), the contents of which are hereby incorporated by reference.

Exemplary of the heterocyclic histamine-1 (H-1) receptor antagonists which are particularly suitable for the present invention, by virtue of their potency in inhibiting neuronally-induced mast cell secretion (as distinct from immunologic degranulation of mast cells), are Azelastine, Azatadine, Cetirizine, Hydroxyzine, Ketotifen, Loratadine and Oxatomide. A particularly suitable agent of this type is Ketotifen (Sigma Chemicals; St. Louis, Missouri) and the pharmaceutically acceptable, non-toxic salts thereof.

Pursuant to the present invention, suitable H-1 receptor antagonists are those that display an affinity for the H-1 receptor that is greater than about 5×10^{-5} M. It is important in this regard to distinguish such high-affinity H-1 receptor antagonists from others of lower affinity which actually cause mast cell degranulation and, hence, should be avoided. See Pini et al., *Agents & Actions* 8: 491-96 (1978). Examples of the latter type of compound appear in Table 1.

Table 1

HISTAMINE-1 RECEPTOR ANTAGONISTS
THAT CAUSE MAST CELL DEGRANULATION

25	<u>Compound</u>	ED ₂₀ for Histamine Release
	Clorphenyramine	10 ⁻³ M
	Cyclizine	5 x 10 ⁻⁴ M
	Diphenhydramine	10 ⁻³ M
30	Promethazine	10 ⁻⁴ M

ED₂₀ = effective dose causing 20% secretion.
M = molar concentration.

Illustrative histamine-3 receptor agonists include N^α-methyl histamine (Calbiochem Corporation, LaJolla, California), N^α-methyl-α(dimethyl) histamine and α,β-difluoro-N^α(fluoromethyl) histamine.

Particularly preferred among the anti-female sex hormones are tamoxifen or NOLVADEX (Zeneca Pharmaceuticals; Wilmington, Delaware), clomiphene or SEROPHENE (Serono Labs; Norwell, Massachusetts), and analogues of gonadotropin releasing hormone (GnRH), a/k/a "leutinizing hormone releasing hormone" (LHRH), such as LUPRON (TAP Pharmaceuticals, Deerfield, Illinois).

Some of the classes of drugs which fall into the above-discussed category of mast cell degranulation-blocking agents are listed below, along with specific examples under each class:

Anthranilic Acid Derivatives

N-(3,4-dimethoxycinnamoyl) anthranilic acids

Anti-Female Sex Hormones

15 Clomiphene
 Mifepristone
 Tamoxifen
 GnRH analogues

Arachidonic Acid Metabolites

20 Leukotriene D₄
 Lipoxin B
 Prostaglandin E₁
 Prostaglandin E₂

Histamine-3 Receptor Agonists

25 N^α-methyl histamine
 N^α-dimethyl histamine
 R^α-methyl histamine
 α,β-dimethyl-N^α(dimethyl) histamine
 α,β-difluoro-N^α(dimethyl) histamine
30 α-methyl,β-fluoro-N^α(fluoromethyl) histamine
 α,β-difluoro-N^α(fluoromethyl) histamine

**Histamine-1 Receptor Antagonists
Characterized by a Three-Ring Structure**Piperidine derivatives

- 5 Azelastine
 Azatadine
 Burfroline
 Doxantrozole
 Forskolin
 Ketotifen
10 Lodoxamide tromethamine
 Loperamide
 Loratadine
 Myricetin
 Oxatomide
15 Pizotifen
 Proxicromil

Piperazine derivatives

- 1-(5-isoquinolinylsulfonyl)-2-methylpiperazine
 1-(1-hydroxy-5-isoquinolinylsulfonyl)piperazine
20 1-[2-hydroxy-3-[(4-hydroxy-3 nitrocoumarin-7-yl)
 oxyl] propyl]-4-(4-chlorobenzyl) piperazine
 Cetirizine
 6-[3-[4-(4-chlorobenzyl)-1-piperazinyl]
 propoxy][1]benzo-pyranc [2,3-d]-1,2,3-triazol-
25 9(1H)-one
 Etodroxizine
 Hydroxyzine
 N-(optionally substituted benzhydryl)-N'-(optionally
 substituted)piperazines
30 9-chloro-5-oxo-7-(1H-tetrazol-5yl)-5H-[1]-
 benzopyrano [2,3-b]pyridine sodium salt
 pentahydrate

Peptides

- 35 L-Asp-Ser-Asp-Pro-Arg
 Cyclosporin A
 Forssman antibody
 Interleukin-1 receptor antagonist
 Lymphocyte Inhibitory Factor (LIF)
 Substance P-receptor antagonist
40 Somatostatin

Phosphatase Inhibitors

- 45 Okadaic acid
 Calyculin
 Quercetin
 Kaempferol
 Nedocromil
 Minocromil
 Cromolyn

Polyamines

Putrescine
Spermidine
Spermine

5 Protein Kinase Inhibitors

- 3-10-Dihydroxy-10[(dimethylamino) methyl]-
2,3,9,10,11,12-hexahydro-9-methyl-9,12,-epoxy-
1H-diindolo[1,2,3-fg-3',2',1'-Kd] pyrrolo [3,4-
c] [2,6] benzodiazocin-1-one (UCH-01)
10 Staurosporin

Proteoglycans

Heparin
Herudin

Quinoline Derivatives

- 15 1,3-Oxazolo[4,5-h]quinolines
2-Carboxypyrimidoquinolines
3-Aminoquinolines

Thiophene Derivatives

- 20 1-Methyl-2(1,3,4-oxadiazol-2(3H)-one-5-yl)
Benzimidazole

Miscellaneous Compounds

- 2-Substituted 3-dimethylamino-5,6-
methylenedioxydines
2-o-Propoxyphenyl-8-azapurin-6-one
25 2-Ethoxyethyl-5-chloro-benzoxazole-2-carboxylate
5-Amino-4-imidazolecarboxamide riboside
7-Methyl-5-propyl-s-triazole[1,5c]pyrimidine-2-amine
7-(2-Hydroxyethoxy)-9-oxoxanthene-2-carboxylic acid
Amiloride
30 Cloxacepride
Deoxycoformycin
Fisetin
Flufenamic acid
Nilidrine hydrochloride
35 p-Bromophenacyl bromide
Pimozide

Phosphatase inhibitors make up another class of
compounds which is characterized by mast cell

degranulation-inhibitory activity. Exemplary compounds in this class are Nedocromil, Minocromil, Kaempferol, Quercetin, 2'-carboxyl-atochromone-5'-yl-2-hydroxypropane derivatives, 1,3-bis [2'-[[acetoxymethyl)oxyl] carbonyl]chromone-5'-yl] 2-hydroxypropane, pyrano [3,2,-g] quinoline-2,8-dicarboxylic acid derivatives, and 5-chlorobenzoxazole-2-carboxylic acid. These compounds and others in this class obstruct a phosphatase from mediating protein dephosphorylation, an integral step in various phosphorylation/dephosphorylation events which are essential to intracellular regulation.

In particular, protein phosphatase inhibitors have been shown to attenuate the release of histamine and leukotriene mediators from human lung mast cells. Peachell et al., *J. Immunol.* 151: 3808 (1993). Phosphatase inhibitors are distinguishable by an ability to inhibit one of the four major classes of phosphatases. Type 1 phosphatase (PP1) dephosphorylates the beta subunit of phosphorylase kinase and is inhibited by thermostable inhibitors 1 and 2 and inactivated by okadaic acid (LC Services Corporation, Nottingham, UK). Type 2 phosphatases include three subtypes (PP2A, PP2B, and PP2C), each of which dephosphorylates the alpha subunit of phosphorylase kinase. Each of PP2A and PP2B are inhibited by okadaic acid. PP2C is insensitive to okadaic acid but is completely dependent on Mg^{2+} .

Accordingly, the class of protein phosphatase inhibitors also includes okadaic acid, analogues of okadaic acid, okadaol and nor-okadaone (LC Services Corporation, *supra*; Biomol, Plymouth Meeting, PA), and okadaic acid methyl ester [see Nishiwaki, *Carcinogenesis* 11: 1837 (1990)], as well as calyculin (LC Services Corporation).

It is not necessary to the operability of the present invention that each member of the foregoing classes inhibits mast cell degranulation to the same extent. The ability of a particular substance to inhibit mast cell degranulation can be readily determined empirically.

Thus, mast cells from rats (pieces of small intestine) or humans (intestinal biopsies) can be obtained; otherwise, RBL cells or immortalized human mucosal-like mast cells (HMC-1) are kept in culture. Any of these cells then are
5 exposed to various concentrations of a putative inhibitory compound, to allow for action by the putative inhibitor at critical sites. Efficacy with respect to inhibition of mast cell degranulation then is ascertained; that is, the intestinal biopsies, explants
10 or cultured cells thus treated are used as such or washed free of unbound test compound before exposure to a stimulus known to induce mast cell degranulation. A compound suitable for use in accordance with the present invention is recognized as one that prevents the stimulus
15 from causing degranulation, presumably by binding to specific receptors or by inactivating a crucial step in the degranulation process.

While it is possible in principle that a particular compound in one of the above-mentioned classes, by virtue
20 of its activity, could be employed in the described manner, it is true generally that the advisability of using any mast cell-degranulation inhibitor according to the present invention must be tempered by clinical considerations, including toxicity, bioavailability,
25 distribution, half-life and clearance, which are conventional to this field. See, for example, GOODMAN AND GILMAN'S THE PHARMACOLOGICAL BASIS OF THERAPEUTICS §§ I & XVII (7th ed.).

Thus, dosage forms for the delivery of the selected
30 mast cell degranulation blocking agent should be those that achieve and retain high levels of the drug in the gastrointestinal tract with reasonable rapidity. These dosage forms are adapted for a variety of conventional routes of administration, including oral, parenteral and
35 rectal.

Pursuant to the present invention, a mast cell degranulation-blocking agent is administered in a pharmaceutically effective amount. The dosage range for

pharmaceutical effectiveness can be determined by reference to standard laboratory tests for inhibition of gastrointestinal mast cell degranulation, as discussed above.

5 A pharmaceutically effective dose for a given mast cell degranulation-blocking agent can be approximated by its ID₂₀, which is the dose necessary to cause 20% inhibition in mast cells when applied to mast cells stimulated *in vitro*. The ID₂₀ values for selected
10 inhibitor compounds are shown in Table 2.

Table 2

COMPOUNDS WHICH INHIBIT RAT MUCOSAL
MAST CELL DEGRANULATION *IN VITRO*

15	Compound	ID ₂₀ for histamine release
	Azatadine	10 ⁻⁷ M
	Hydroxyzine	5 x 10 ⁻⁵ M
	Ketotifen	10 ⁻⁵ M
	N ^α -methyl histamine	10 ⁻⁷ M
20	Oxatomide	10 ⁻⁷ M
	Spermine	5 x 10 ⁻⁵ M
	Tamoxifen	5 x 10 ⁻⁵ M

25 ID₂₀ = inhibitory dose causing 20% inhibition.
M = molar concentration.

The agent can best be administered systemically or directly in the intestinal loops (suppository, enema or injection) of the live mammal. The mast cell inhibitory effect of certain of these compounds has been described
30 for non-intestinal mast cells. See, for example, Vliagoftis et al., *Biochem. Pharmacol.* 43: 2237-45 (1992), Rozniecki et al. in *PROC. 8TH INT'L CONGRESS IMMUNOLOGY* 159 (Budapest 1992), Rozniecki et al., *FASEB J.* 6: A1554 (1992), and Rachmilewitz et al.,
35 *Gastroenterol.* 102: A235 (1992).

An assay by which efficacy of a mast cell degranulation-blocking agent can be tested systemically in the present context is the rat-immobilization assay, in which a rat is pre-treated with an agent and then

restrained for one hour in a clear plexiglass cylinder. After the immobilization, a specific gastrointestinal mast cell mediator, rat mast cell protease II (RMCPII), is measured. This assay is accepted as a non-ulcerogenic model of stress. See Williams et al., *Am. J. Physiol.* 253: 582-86 (1987).

Results from testing ketotifen in the rat-immobilization assay are presented in Table 3 below. Ketotifen was administered intraperitoneally to rats (1 mg/kg body weight, twice daily) for four days prior to immobilization.

Table 3
INHIBITION BY KETOTIFEN OF
RAT INTESTINAL RMCPII IN A STRESS MODEL

Time (min)	Immobilized (n = 3)	Non-immobilized (n = 3)	Pre-treated w/ ketotifen (n = 3)
60	4.85 [*]	0.3	0.5
120	12.4 [*]	0.7	0.8 ^{**}

^{*}ng/ml; n = number of rats tested

* p < 0.05 vs. non-immobilized

** p < 0.001 vs. immobilized

Table 3 shows that intestinal mast cells are activated during stress to secrete a specific proteolytic enzyme, RMCPII, and that ketotifen acts as a mast cell degranulation-blocking agent in preventing secretion of RMCPII from such cells. The equivalent proteolytic enzyme in humans is tryptase, which also can be measured from colonic fluid or biopsies.

It is essential to the present invention that the selected agent inhibit mucosal mast-cell secretion. Thus, mast cell degranulation-blocking agents, such as disodium cromoglycate (CROMOLYN), which do not have an inhibitory effect specifically on mucosal mast cells, see Pearce et al., *J. Immunol.* 128: 2481-86 (1982), are not within the scope of the present invention. No work has

so far involved H3 receptor agonists in the inhibition of gastrointestinal mast cell degranulation.

5 A mast cell degranulation-blocking agent can be administered, according to the present invention, in the form of the active agent itself or as a pharmaceutically acceptable salt of the active agent. The phrase "pharmaceutically acceptable salt" denotes a non-toxic, substantially non-irritating salt of the compound used. Typical salts are ammonium salts and salts containing an
10 cation that is an alkali metal or alkaline earth metal, such as sodium, potassium, calcium or magnesium. The category of suitable cations for the salt also includes sulphate, phosphate, tartrate and citrate. Other acceptable salts are those with non-toxic organic acids,
15 such as fatty acids of one to six carbon atoms in length.

Pursuant to the present invention, a mast cell degranulation-blocking agent can be delivered via methods known for drug administration. The agents are typically administered as pharmaceutical compositions in
20 combination with pharmaceutically acceptable carriers. Such compositions may be prepared from conventional materials by procedures well known in the art.

Compositions within the present invention can be adapted for oral or parenteral administration, as well as
25 for enteral administration or through mucous membranes, such as transdermally. An oral formulation which will keep the drug for an extended time in the gastrointestinal tract is preferred, since such a formulation allows long exposure of the mucosal mast
30 cells present there to the drug.

Forms suitable for oral administration include tablets, dispersible powders, granules, capsules, syrups, elixirs and suspensions. Also preferred in this regard are encapsulated forms. For instance, a preferred mode
35 of administration would be via liposomes, multilamellar phospholipid vesicles, or lipospheres, which protect the active agent from destruction in the stomach and allows for slow release in the intestine as each phospholipid

layer is dissolved, resulting in sustained high intestinal levels of the drug. Alving, C.R., S. Shichijo, I. Mattsby-Baltzec, R.L. Richards and N.M. Wassef, "Preparation and Use of Liposomes in Immunologic Studies," in *LIPOSOME TECHNOLOGY*, 2nd ed., vol. III, G. Gregoriadis, ed., CRC Press, Boca Raton, FL, pp.317-343, 1984; Amselem, S., C.R. Alving, and A.J. Domb, *Polymers for Advanced Technologies*, 3:351-357 (1992); and Amselem, et al. *Vaccine Research* 1: 383-385 (1992).

Compositions for oral use contain one or more conventional adjuvants, such as sweetening agents, flavoring agents, coloring agents and preserving agents, in order to provide a presentable and palatable preparation. Tablets may contain the active ingredients in a mixture with conventional pharmaceutically acceptable excipients. These include inert carriers, such as calcium carbonate, sodium carbonate, lactose, and talc; granulating and disintegrating agents, such as starch and alginic acid; binding agents such as starch, gelatin acacia; and lubricating agents, such as magnesium stearate, stearic acid and talc. Tablets may be uncoated or coated by known techniques to delay disintegration and absorption in the gastrointestinal tract, thereby providing a sustained action over a longer period of time. Similarly, suspensions, syrups and elixirs may contain active ingredients in mixture with any of the conventional excipients utilized in the preparation of such compositions. This includes suspending agents such as methylcellulose, tragacanth and sodium alginate; wetting agents such as lecithin, polyoxyethylene stearate or polyoxyethylene sorbitan monolaurate; and preservatives. Capsules may contain the active ingredients alone or in an admixture with an inert solid carrier, such as calcium carbonate, calcium phosphate or kaolin. These pharmaceutical compositions may contain up to 90% of active ingredients in combination with the carrier or adjuvant. The compounds preferably are put up in unit dosage forms, particularly for oral administration, for

example, in 20 mg tablets. Such forms may contain the active ingredient separately, for example in separate layers.

5 The agents can be administered in sustained release form or in divided dosages.

Oral administration often is preferred if the mast cell degranulation-blocking agent is orally active. Transdermal administration also is preferred. More specifically, in the case of mast cell degranulation-
10 blocking agents which are substantially destroyed or deactivated upon oral administration, or where a more prolonged duration of activity is desired, a mast cell degranulation-blocking agent also can be administered transdermally or via other body membranes, such as
15 rectally.

A preferred formulation within the present invention comprises both a suitable mast cell-degranulation inhibitor and an agent that inhibits or counteracts intestinal motility, thereby increasing the transit time
20 and, hence, the effect of the inhibitor in the intestinal tract. Such a suitable antimotility agent is dicyclomine capsules, such as BENTYL (Taylor Pharmaceutical Company, Decatur, Illinois). The use of an antimotility agent in this context is especially preferred when the mast cell-
25 degranulation inhibitor is contained in a lipid-containing delivery vehicle, as discussed above. Thus, the antimotility agent can be "co-administered" with the delivery vehicle, i.e., can be administered in such a way that the motility-counteractive influence of the agent
30 increases the residence time of the delivery vehicle in the intestine.

The following illustrative examples further describe the present invention.

Example 1

35 The following experiment was performed in order to evaluate rat mucosal mast cell (MMC) degranulation by

measuring rat mast cell protease II (RMCP II), a specific marker for MMC, and by electron microscopy.

Rats were treated intragastrically for four (4) days with either 0.5 ml of saline containing ketotifen (1 mg/kg twice daily) or saline alone. On day five, the animals were euthanized, ileum was removed and ileal explants were placed in organ culture. The explants were incubated at 37°C in a medium containing either 5 µg/ml of toxin A in 0.01 ml of 50 mM Tris buffer, or buffer alone. Aliquots (50 µl) of medium were collected at specified times and RMCP II levels were determined. Results are expressed in Table 4 as nanograms of RMCP II released per milligram of tissue wet weight (Pothoulakis et al., *Gastroenterology* 105:701-707, 1993).

Table 4

INHIBITION BY KETOTIFEN OF RAT INTESTINAL
RMCP II SECRETION INDUCED BY TOXIN A

Time (min)	Buffer (n = 10)	Toxin A (n = 7)	Ketotifen ⁺ Toxin A (n = 6)
30	3.4 ± 0.8	5.0 ± 1.2	2.9 ± 1.0
60	6.2 ± 0.9	7.2 ± 1.7	3.0 ± 1.1 ⁺
120	7.6 ± 1.2	12.6 ± 2.1 [*]	4.2 ± 0.9 ⁺⁺
180	11.3 ± 2.1	24.0 ± 4.8 ^{**}	6.7 ± 1.9 ⁺⁺

* p < 0.05 vs buffer

** p < 0.01 " "

+ p < 0.05 vs Toxin A

++ p < 0.01 " "

Electron microscopy showed significant mucosal mast cell degranulation in rat ileum within 15 minutes of exposure to toxin A, and complete degranulation at 60 minutes. These results, obtained by J. Castagliuolo, J.T. LaMont, R. Letourneau, C. Kelly, J.C. O'Keane, A. Jaffer, T.C. Theoharides and C. Pothoulakis, demonstrate that ketotifen effectively inhibits the release of RMCP II from rat ileal explants and, hence, inhibits mucosal mast cell degranulation.

Similar results were obtained using the substance P receptor antagonist (Pothoulakis et al., Proc. Natl. Acad. Sci., USA 91:947-951, 1994).

Example 2

5 A female patient, age 49, suffered from irritable
bowel syndrome for approximately 10 years. Symptoms were
so severe that she had difficulty in functioning as a
nurse, mother and wife. After treatment with hydroxyzine
10 hydrochloride (ATARAX; 50 mg/day) for three months, her
gastrointestinal discomfort was almost completely
eliminated, and she was able to resume her professional
and family responsibilities. On several occasions when
she attempted to withdraw from the use of hydroxyzine
hydrochloride, she experienced a marked increase in
15 symptoms of irritable bowel syndrome. The symptoms
disappeared when she resumed taking the medication.

Example 3

20 A female patient, age 32, had a four-year history of
irritable bowel syndrome and had difficulty in carrying
out her professional duties as a nurse. She was treated
for 4 months with 50 mg/day of hydroxyzine pamoate
(VISTARIL). She experienced a noticeable improvement of
her symptoms and was able to resume her professional
activities.

25 Example 4

A female patient, age 39, suffered from irritable
bowel syndrome for over two years. Her symptoms included
constant pain at the connection of the buttocks with the
upper legs, as well as painful defecation. After
30 treatment with ketotifen (ZADITEN; 4 mg/day) for two
months, her pain subsided significantly.

Example 5

A female patient, age 39, with a five-year history
of irritable bowel syndrome, experienced constant

abdominal pain which was not relieved by any of a variety of narcotic and non-narcotic analgesics and anti-inflammatory drugs. Her symptoms were exacerbated during menstruation. Treatment with tamoxifen citrate (NOVADEX;
5 40 mg/day, administered orally) for two months substantially eliminated her pain.

Example 6

A female patient, age 32, with a two-year history of abdominal migraine, experienced constant gastrointestinal
10 pain which was considerably worse immediately premenstrually. After treatment with the Gn-RH analogue LUPRON (3 µg/day subcutaneous injection) for four months, she was symptom-free.

WHAT IS CLAIMED IS:

1. A method of manufacturing a medicament for the treatment of an endogenous, painful gastrointestinal condition of non-inflammatory, non-ulcerative origin which comprises a mast cell degranulation-blocking agent in an amount sufficient to affect said condition in a pharmaceutically acceptable carrier.

2. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is a naturally occurring polyamine.

3. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is selected from the group consisting of spermine, spermidine, and metabolites thereof.

4. The composition according to claim 3, wherein said mast cell degranulation-blocking agent is spermine.

5. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is administered orally.

6. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is contained in a lipid-containing delivery vehicle.

7. The composition according to claim 6, wherein said lipid-containing delivery vehicle is co-administered with an antimotility agent.

8. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is a heterocyclic histamine-1 receptor antagonist.

9. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is selected from the group consisting of azatadine, azelastine, cetirizine, forskolin, hydroxyzine, ketotifen and oxatomide.

10. The composition according to claim 9, wherein said mast cell degranulation-blocking agent is ketotifen.

11. The composition according to claim 10, wherein said ketotifen is in the form of a pharmaceutically acceptable salt.

12. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is a histamine-3 receptor agonist.

13. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is selected from the group consisting of N^α-dimethyl histamine, R^α-methyl histamine, α,β-dimethyl-N^α(dimethyl) histamine, α,β-difluoro-N^α(dimethyl) histamine, α-methyl,β-fluoro-N^α(fluoromethyl) histamine and α,β-difluoro-N^α(fluoromethyl) histamine.

14. The composition according to claim 13, wherein said mast cell degranulation-blocking agent is α,β-difluoro-N^α(fluoromethyl) histamine.

15. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is an anti-female sex hormone.

16. The composition according to claim 1, wherein said mast cell degranulation-blocking agent is clomiphene, mifepristone, tamoxifen and a GnRH analogue.

17. The composition according to claim 16, wherein said mast cell degranulation-blocking agent is tamoxifen.

18. The composition according to claim 16, wherein said mast cell degranulation-blocking agent is LUPRON.

19. The composition according to claim 18, wherein said LUPRON is administered subcutaneously.

20. The composition according to claim 1, wherein said condition is abdominal migraine.

21. The composition according to claim 1, wherein said condition is irritable bowel syndrome.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/13, 31/445, 31/55, 31/495, 31/35, 31/415, 31/135, 31/575, 38/09	A3	(11) International Publication Number: WO 95/21611 (43) International Publication Date: 17 August 1995 (17.08.95)
(21) International Application Number: PCT/US95/01392 (22) International Filing Date: 9 February 1995 (09.02.95) (30) Priority Data: 08/193,597 9 February 1994 (09.02.94) US (71) Applicant: KOS PHARMACEUTICALS, INC. [US/US]; Suite 2502, 1001 South Bayshore Drive, Miami, FL 33131 (US). (72) Inventor: THEOHARIDES, Theoharis, C.; 14 Parkman Street, Brookline, MA 02146 (US). (74) Agents: MELOY, Sybil et al.; Foley & Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 5 October 1995 (05.10.95)	
(54) Title: COMPOSITIONS COMPRISING A MAST CELL DEGRANULATION-BLOCKING AGENT FOR TREATMENT OF ENDOGENOUS, PAINFUL GASTROINTESTINAL CONDITIONS OF NON-INFLAMMATORY, NON-ULCERATIVE ORIGIN (57) Abstract A method for treating endogenous, painful gastrointestinal conditions of non-inflammatory, non-ulcerative origin, such as abdominal migraine and irritable bowel syndrome, entails administering a pharmacologically effective amount of a mast cell degranulation-blocking agent, e.g. a naturally occurring polyamine, a heterocyclic histamine-1 receptor antagonist, a histamine-3 receptor agonist or an anti-female sex hormone.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/US 95/01392

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/13 A61K31/445 A61K31/55 A61K31/495 A61K31/35
A61K31/415 A61K31/135 A61K31/575 A61K38/09

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DIG DIS SCI (UNITED STATES), SEP 1993, VOL. 38, NO. 9, PAGE(S) 1590-5, Weston AP et al 'Terminal ileal mucosal mast cells in irritable bowel syndrome.'	1,5,21
Y	see the whole document	1-7,20, 21
X	SCAND J GASTROENTEROL SUPPL (NORWAY), 1987, VOL. 130, PAGE(S) 7-13, Read NW 'Irritable bowel syndrome (IBS)--definition and pathophysiology.'	1,21
Y	see the whole document	1-7,20, 21
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- '&' document member of the same patent family

Date of the actual completion of the international search

31 May 1995

Date of mailing of the international search report

11.09.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Stierman, B

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/US 95/01392

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AM. J. GASTROENTEROL., vol.87, 1992 pages 55 - 57 G.F. STEFANINI 'Oral disodium cromoglycate treatment on irritable bowel syndrome: An open study on 101 subjects with diarrheic type'	1,5,21
Y	see the whole document	1-7,20, 21
Y	GLAS SRPSKE AKADEMIJE NAUKA, vol.24, 1971 pages 43 - 50 P. STERN ET AL. 'Contribution to pharmacology of spermidine' see abstract	1-7,20, 21
Y	CLIN. EXP. PHARMACOL. PHYSIOL. (ENGLAND), 1986, VOL. 13, NO. 1, PAGE(S) 87-90, 'Comparison of the inhibitory effects of spermine, papaverine and adrenaline upon isolated segments of the rat small intestine' see the whole document	1-7,20, 21
Y	70TH ANNUAL MEETING OF THE FEDERATION OF AMERICAN SOCIETIES FOR EXPERIMENTAL BIOLOGY, ST. LOUIS, MO., USA, APR. 13-18, 1986.;& FED PROC., VOL. 45, NO. 3, PAGE(S) 437, 1986. CARTAGENA A ET AL 'INHIBITION OF MAST CELL DEGRANULATION BY NATURAL POLYAMINES' see abstract	1-7,20, 21
Y	J CLIN GASTROENTEROL (UNITED STATES), JUN 1981, VOL. 3, NO. 2, PAGE(S) 153-6, PAGE JG ET AL 'Treatment of the irritable bowel syndrome with Bentyl (dicyclomine hydrochloride).' see the whole document	7
Y	WO,A,93 12773 (KOS PHARM., INC.) 8 July 1993 see the whole document	1-7,20, 21
Y	GASTROENTEROLOGY,, VOL. 99, NO. 6, PAGE(S) 1668-1674, 1990. AURICCHIO S ET AL 'AMINES PROTECT IN-VITRO THE CELIAC SMALL INTESTINE FROM THE DAMAGING ACTIVITY OF GLIADIN PEPTIDES' see the whole document	1-7,20, 21

-/--

INTERNATIONAL SEARCH REPORT

Intern al Application No

PCT/US 95/01392

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US,A,4 507 321 (I.H. RAISFELD) 26 March 1985 see column 2 see column 8 ---	1-7,20, 21
A	BIOCHEM. PHARMACOL. (ENGLAND), 1983, VOL. 32, NO. 11, PAGE(S) 1733-1736, AIHARA H. ET AL 'Polyamine inhibition of gastric ulceration and secretion in rats' see the whole document ---	1-7,20, 21
A	AM. J. PHYSIOL., 1993, VOL. 264 pages G367 - G374 MCCORMACK, SHIRLEY A. ET AL 'Polyamines are necessary for cell migration by a small intestinal crypt cell line.' see the whole document -----	1-7,20, 21

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 95/01392

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
please see enclosure!
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

please see enclosure!

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

2-4, 1,5-7,20,21 (part.)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

LACK OF UNITY OF INVENTION

1. Use of a naturally occurring polyamine in the manufacture of a medicament for the treatment of an endogenous, painful gastrointestinal condition of non-inflammatory, non-ulcerative origin.
(Claims 2-4 completely; claims 1,5-7,20,21 partially).
2. Use of a heterocyclic histamine-1 receptor antagonist in the manufacture of a medicament... (see above).
(Claims 8-11 completely; claims 1,5-7,20,21 partially).
3. Use of a histamine-3 receptor agonist in the manufacture ... (see subject 1).
(Claims 12-14 completely; claims 1,5-7,20,21 partially).
4. Use of an anti-female sex hormone in the manufacture ... (see subject 1).
(Claims 15-19 completely; claims 1,5-7,20,21 partially).

Please note that a possible further search for the subjects not searched yet might result in further objections a posteriori concerning unity.

The problem to be solved by the present application is to provide a treatment for an endogenous, painful gastrointestinal condition of non-inflammatory, non-ulcerative origin (especially abdominal migraine and irritable bowel syndrome).

The solution proposed in the present application is to use a mast cell degranulation-blocking agent as the active therapeutic substance.

The common technical feature and therefore the linking inventive concept, which, a priori, unifies the different groups of compounds, is the use of a mast cell degranulation-blocking agent as the therapeutic agent.

From Dig. Din. Sci., 1993, Vol. 38, No. 9, pp. 7-13, it is known that mast cell stabilizers like e.g. Cromolyn, Ketotifen, H1-receptor antagonists etc. are useful in the treatment of irritable bowel syndrome.

In Scand. J. Gastroenterol., 1987, Vol. 130, Suppl., pp. 7-13, the use of mast cell stabilizers in the treatment of irritable bowel syndrome is described.

./.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US95/01392

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

From Am. J. Gastroenterol., 1992, Vol. 87, pp. 55-57, it is known, that the mast cell stabilizers Disodium Cromoglycate and Doxantrazole are useful against irritable bowel syndrome.

It is apparent from these documents that the proposed solution of the problem, namely the use of a mast cell degranulation-blocking agent as the active therapeutic substance against the claimed gastrointestinal disorders, is not novel.

In the absence of a special technical feature linking the different groups of compounds used, which, as the man skilled in the art knows, differ substantially in their mode of action as well as in their structure, the application lacks unity of invention and must be divided a posteriori according to the different modes of action of the different groups of compounds.

In the present application no further technical feature(s) can be distinguished that can be regarded as a "special technical feature" involved in the technical relationship among the different inventions.

Consequently, the present application lacks unity of invention, and the different solutions not belonging to a common inventive concept are identified as different subjects listed in the communication pursuant to Art. 17(3)(a) PCT.

Each of the inventions listed is a distinct invention, characterized by its own special technical feature, defining the contribution which each of the claimed inventions, considered as a whole, makes over the prior art.

Searching these different subjects would have caused major additional searching efforts.
Only the first subject was searched.

Expressions like "naturally occurring polyamine", "antimotility agent", "heterocyclic histamine-1 receptor antagonist", "histamine-3 receptor agonist", "anti-female sex hormone", "GhRH analogue" etc. do not make sufficiently clear, which specific compounds are meant.

A compound is not sufficiently described as "metabolite" of another compound.

In view of this the search had to be/would have to be restricted to the compounds explicitly mentioned in the claims and to the general inventive concept.

The description of the disorder in claim 1 does not make completely clear, which diseases are meant.

The search had to be/would have to be restricted to the diseases explicitly mentioned in the claims and to the general inventive concept.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US95/01392

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

Incomplete search

Claims searchable incompletely: 1-21

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 95/01392

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9312773	08-07-93	JS-A- 5250529 AU-B- 3422793 EP-A- 0618796	05-10-93 28-07-93 12-10-94
US-A-4507321	26-03-85	NONE	